

Overcoming the disconnect between interaction networks and biodiversity conservation and management

Gabriel Dansereau^{1,2} João Braga³ Gentile Francesco Ficetola^{4,5} Núria Galiana⁶ Dominique Gravel^{2,7}
Luigi Maiorano⁸ José M. Montoya⁹ Louise O'Connor¹⁰ Laura Pollock¹¹ Wilfried Thuiller⁵
Timothée Poisot^{1,2} Ceres Barros^{12,13,14}

¹ Département de sciences biologiques, Université de Montréal, Montréal, Québec, Canada

² Québec Centre for Biodiversity Science, Montréal, Québec, Canada

³ Ecofish Research Ltd., Courtenay, British Columbia, Canada

⁴ Department of Environmental Science and Policy, Università degli Studi di Milano, Milano, Italy

⁵ Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LECA, F-38000 Grenoble, France

⁶ Department of Biogeography and Global Change, National Museum of Natural Sciences, Madrid, Spain

⁷ Département de biologie, Université de Sherbrooke, Sherbrooke, Québec, Canada

⁸ Department of Biology and Biotechnologies “Charles Darwin”, Sapienza University of Rome, Italy

⁹ Theoretical and Experimental Ecology Station, CNRS, Moulis, France

¹⁰ Biodiversity, Ecology and Conservation Group, International Institute of Applied Systems Analysis, Laxenburg, Austria

¹¹ Biology Department, McGill University, Montréal, Québec, Canada

¹² Pacific Forestry Centre, Canadian Forest Service, Natural Resources Canada, Victoria, British Columbia, Canada

¹³ Department of Forest Resources Management, University of British Columbia, Vancouver, British Columbia, Canada

¹⁴ Département des sciences du bois et de la forêt, Université Laval, Québec City, Québec, Canada

Correspondence to:

Gabriel Dansereau — gabriel.dansereau@umontreal.ca

Decision-makers need to act now to halt biodiversity loss, and ecologists must provide them with relevant species interaction indicators to inform on community- and ecosystem-level changes. Yet, the integration of ecological networks into conservation is still virtually nonexistent. Here, we discuss challenges and

opportunities related to uncertainty, interpretability and relevance of network metrics applied to conservation. We argue that existing data and methodologies are sufficient to generate network information usable for conservation, and to overcome existing challenges. Interaction network indicators must meet criteria important to decision-makers and be tied to specific conservation goals, which requires academics to better engage with practitioners. We suggest network robustness as an indicator for biodiversity management and showcase it in a workflow to inform decision-making.

This work is released by its authors under a CC-BY 4.0 license



Last revision: *December 6, 2024*

Highlights

- Practitioners and scientists increasingly need multi-species and whole-ecosystem indicators that allow integrating species interaction networks into biodiversity conservation and management.
- Explicit and quantitative integration of ecological network indicators into conservation is still lacking due to challenges with network uncertainty and accessibility to practitioners.
- The resulting gap between network science and management leads to decisions being made without considering available scientific knowledge.
- We identify opportunities in closing this gap. Despite uncertainty, the field of network ecology is mature enough to offer quantitative insights into ecosystem responses to environmental changes.
- Simple network metrics that fit criteria important to decision-makers and can be used with current data and models are promising starting indicators to inform conservation and management.

1

2 **Can interaction network knowledge be quantitatively used for biodiversity** 3 **conservation and management?**

4 The need to shift from single-species conservation approaches to multi-species and whole ecosystem
5 approaches has long been recognized [1,2]. Network information can provide a new perspective for whole
6 ecosystem assessments in biodiversity conservation and management. Preserving species interactions can
7 ensure long-term population persistence and maintain ecosystem functions and services [3,4]. Focusing on
8 ecological networks as conservation targets promotes the stability of populations and ecosystem functions and
9 minimises negative outcomes regarding species extinctions [5–7]. Recent reviews list specific interaction
10 network metrics that decision-makers can use [8]. Implicit network information has already been integrated into
11 conservation planning, for example through consideration of keystone species with disproportionate effects on
12 their communities, which should facilitate the uptake of network-based biodiversity indicators in
13 decision-making [2,9,10, see Box 1].

14 Despite the potential benefits, conservation practices rarely explicitly consider information derived from
15 measures of the structure of ecological networks. Conservation policy and practice still heavily focus on single
16 species and habitats. Uncertainty about network structure and responses to human disturbances mirrors

17 concerns in macro-ecological and ecosystem models [11,12]. Additionally, identifying which interaction
18 network metrics are suitable biodiversity indicators with clear interpretation for conservation remains
19 challenging.

20 Decision- and policy-makers must act now to bend the curve of extinction and accelerate ecosystem recovery
21 [13,14]. Ecologists need to provide them with useful network and ecosystem-wide information. For instance,
22 protected area planning could prioritise regions where mutualistic interaction partners or prey and predators
23 overlap [15], or where there is high trophic diversity and redundancy, enhancing robustness to extinctions [16].
24 Moreover, since interaction network structure is linked to ecosystem functioning and ecosystem service
25 provision, focusing on network metrics changes for conservation targets should ensure ecosystem stability and
26 service delivery [e.g., pollination, pest control, food production, 5,7,17]. Given the global goals to maintain
27 ecosystem services [Goal B of the Kunming-Montreal Global Biodiversity Framework, 18], assessing network
28 structure stability changes should help managers and decision-makers prioritise areas to maintain ecosystem
29 functioning and resilience [5,19].

30 Here, we identify the major challenges and opportunities in incorporating interaction network information into
31 biodiversity conservation and ecosystem management. We demonstrate how simple approaches and indicators
32 can provide relevant information for managers. Our focus is on probabilistic and binary species interaction
33 networks, where nodes represent species and links represent the probability or presence of an interaction [20],
34 rather than energy flow networks already covered by Fath et al. [8]. Additionally, we present a perspective
35 where networks are used as biodiversity indicators and, in a forecasting context, to evaluate network responses
36 to future environmental change scenarios and management strategies. Despite challenges relating to
37 uncertainty, interpretability and relevance, we argue that we have sufficient scientific evidence and tools to apply
38 network concepts to management and conservation in the face of global change. In particular, testing and
39 exploring network indicators can accelerate the establishment of operational monitoring frameworks.

Box 1 - Trophic role of keystone species

Explicitly considering networks in conservation and decision-making (i.e. by monitoring and managing for network-derived properties) is not a drastic shift, as networks are often implicitly included in conservation decisions and recovery plans. The keystone species concept, frequently mentioned in conservation literature [e.g., 2,21] and highlighted by initiatives focused on rewilding and ecological restoration [22,23], is linked to the disproportionate effects some species have on their (trophic) networks [24, also see

40

25 for the diverse roles of species identified as keystones]. Similarly, several large carnivores have been associated with trophic cascades, where effects of predator declines propagated across food webs to herbivores, mesopredators, and beyond [26]. This reflects network consideration through species' effects on others, even if network-specific properties are not explicitly quantified – i.e. metrics like connectance, species trophic level, or centrality do not explicitly enter planning or decision-making.

Importantly, keystone species are often tied to quantified conservation targets. For example, prairie dogs (*Cynomys spp.*) are considered keystone species due to their important ecosystem functions and large impact compared to other herbivores, which are not replicated by other species [27,28]. The Recovery Strategy and Action Plan for the Black-tailed Prairie Dog (*Cynomys ludovicianus*) in Canada identifies it as a conservation priority due to its keystone status, crucial for the recovery of the Black-footed Ferret (*Mustela nigripes*) and serving as a vital food source for several other at-risk species [29]. Conservation targets for Black-tailed Prairie Dogs in Canada include maintaining a minimum area of occupancy of 1,400 ha across 20 colonies and a minimum average population density of 7.5 individuals/ha by 2040, ensuring at least an 80% probability of population persistence over 50 years [29].

The implicit consideration of network structure in conservation targets can facilitate the uptake of new network-based indicators by practitioners and decision-makers. Indeed, knowing this structure provides additional ways to identify which species are potential keystones, beyond their emblematic nature [30]. Other forms of network-thinking are similarly part of management considerations, such as spatial ecological networks planning [31] and ecosystem-based management [11]. Explicitly considering network-based indicators will complement these forms of network-thinking and enhance conservation assessments to include ecosystem-wide components.

41

42 **Challenges & opportunities**

43 The explicit integration of network information into management and conservation faces several challenges
44 linked to uncertainties and lack of interpretability and relevance of network metrics for practitioners. These
45 challenges will hinder making effective decisions, for example on what biodiversity and network-related
46 properties need to be measured and monitored, what conservation targets and management actions should be

47 applied, how often to re-evaluate decisions, etc. Hence, we can expect challenges at different stages of
48 management planning and decision-making [e.g. 32], such as the evaluation of current conditions or upon
49 decisions on possible actions (e.g. responsive, preventative, etc.).

50 **Uncertainty**

51 **Network Structure and Composition**

52 There is uncertainty in network structure, composition, and variation across space and time, which affects
53 conservation assessments and actions [33,34]. Empirical studies on networks are often spatially disjointed,
54 biased geographically and depending on interaction types, and rarely replicated [35–37]. Sampling biases can
55 distort reported network patterns [38,39]. Terrestrial and freshwater food webs are less studied than marine
56 ones, often with different research objectives [e.g., determining the effect of environmental factors, rather than
57 investigating management-related elements such as sustainability, 35,40]. Such deficits of information may
58 prove problematic when conservation decisions need to be made.

59 Despite these challenges, existing methodologies can help integrate network information into conservation,
60 while empirical data continue to be gathered. Networks can be constructed from extensive, long-term
61 monitoring datasets to analyse food web structure and temporal stability [41,42]. Building metawebs of all
62 potential interactions in a region or species pool, like the pan-European terrestrial tetrapod metaweb
63 [TETRA-EU, 43], provides an “upper ceiling” for possible interactions [44,45]. Metawebs can inform
64 broad-scale assessments and have already been used to derive spatially explicit network properties and generate
65 conservation-relevant information [46–48]. For instance, Albouy et al. [46] used a metaweb to examine
66 robustness to extinction scenarios for marine food webs, showing higher robustness in coastal waters compared
67 to open waters and highlighting some potential to absorb perturbations. Moreover, metaweb inference
68 approaches allow us to circumvent the lack of available local interaction data [45] and, when used with
69 probabilistic networks, to integrate uncertainty and variation in network structure across space [49]. Network
70 properties and their uncertainties can therefore be measured for broad-scale assessments of variation in network
71 structure, and to derive network indicators that can be used to inform decisions and planning (Boxes 2-3). As
72 new empirical data becomes available, these predictions can be evaluated, refined, and become more
73 informative [50]. We discuss the challenges surrounding their validation in our [Concluding Remarks](#).

74 **Network Responses to Environmental Change**

75 Uncertainty exists in how networks will respond to environmental changes and disturbances, particularly for
76 interaction rewiring and changes in interaction strength. Questions remain on the extent of rewiring due to
77 species turnover versus prey switching and behavioural adaptation, and how these changes will propagate across
78 trophic levels.

79 While data gaps exist, modelling and inference can explore the limits of network rewiring under current or
80 future conditions (Box 3). Rewiring potential is likely captured in existing and inferred metawebs [51], which
81 can be combined with simulations to anticipate network changes. For instance, Dansereau et al.'s [49] approach
82 can be extended to explore climate change impacts on network structure, given the dual uncertainty in species
83 interactions and future species ranges. Moreover, network models (and information) do not need
84 well-constrained or low uncertainty predictions before they can inform management decisions on interventions
85 like species eradication, especially if they tend to correctly identify whether effects on other species will be
86 positive or negative [52]. Model uncertainty can also be high despite high quality data [52]. Regardless of its
87 generality, this result suggests that the performance of a model should be monitored whenever new data are
88 added. Similar trends of model change in performance with additional data have been reported in the study of
89 species distributions [53].

90 Approaches to include specific types of network response uncertainty in conservation and management have
91 also been proposed. Van Kleunen et al. [54] suggested a multi-step framework for decision-making under
92 uncertainty for species introduction into ecological networks, based on conservation decision theory. This
93 framework includes: the identification of management objectives, the evaluation of outcomes for management
94 (including multiple outcomes, evaluation of trade-offs, and assessment of uncertainty), and the improvement of
95 future predictions through an adaptive management framework. Van Kleunen et al.'s [54] decision-making
96 approach can be applied now, despite uncertainties, to guide management of species introductions.

97 **Compounding Uncertainty in Change Types**

98 There is compounding uncertainty in the type and strength of change applied to a network. Climate uncertainty,
99 for instance, results from uncertainty in future greenhouse gases emissions (i.e. emission scenario uncertainty),
100 in climate processes (general circulation model uncertainty) and their stochasticity (model run uncertainty). For
101 networks, we add uncertainty in changes resulting from disturbance regimes (e.g. fire, drought, pests) and in

102 species distribution predictions [which can result from direct impacts of abiotic change, of disturbance regimes
103 and of biotic changes that may be linked to network structure itself, 55,56]. If accounted for simultaneously,
104 these uncertainties will inevitably lead to high variance in predicted network responses.

105 We can estimate some uncertainty through backcasting: past environmental changes are used to predict changes
106 in network metrics that are cross-validated against observed past networks. Fisheries data, for instance, allow
107 reconstructing well-resolved networks over time, which can be related to known environmental changes [57–59]
108 and be used to calibrate predictive network models, like bayesian networks [60]. Backcasting models, used as
109 ex-ante scenarios of change, have been successfully used to simulate and assess the effectiveness of
110 conservation actions on ecosystem services [61].

111 Simulating scenarios of change can also help delimit the possible changes in network structure [Box 3, 62].

112 When combined with metrics of network change and sensitivity to disturbance, these projections can be used to
113 identify target areas that show fragility to an array of scenarios and are of special concern, or that show less
114 fragility and could be considered refugia. They can also highlight problematic or incomplete sampling.

115 Projections will also serve to perform validation and assess indicator behaviour in an empirical setting, whether
116 through existing data or backcasting exercises, which could lead to network-specific monitoring programs.

117 **Interpretability and relevance**

118 Network metrics are often not intuitive or deemed relevant for practitioners and decision-makers. Many metrics
119 are complex and may not clearly correlate with ecosystem- and species-level responses, particularly in applied
120 contexts. For instance, omnivory and network motifs are tied to food web persistence and extinction risks
121 [63,64], highlighting their ecological relevance. On the other hand, while network nestedness indicates a buffer
122 against extinctions and fluctuations in mutualistic networks, this is less clear in antagonistic networks [7].

123 Connectance has also been tied in contrasting ways to network stability [e.g., higher connectance leading to
124 increases or decreases of invasion success rates given invader trophic levels, 65, higher connectance linked to
125 higher robustness to extinction, but larger extinction cascades, 66].

126 Not all network metrics are suitable as conservation indicators, nor do they need to be. Several have been
127 reviewed for their relevance and limitations in achieving conservation goals (Louise O’Connor, PhD thesis,
128 Université Grenoble Alpes, 2022ⁱ; see Table 1 therein). For example, prioritising trophic networks with
129 stabilising motifs when selecting protected areas can help achieve ecological resilience goalsⁱ. This information

130 can already be used towards conservation planning but it needs to be both accepted by and available to
131 decision-makers and managers.

132 First, metrics must meet decision-makers' criteria. The ROARS (being Relevant, Objective, Available,
133 Realistic, Specific) and SMART (Specific, Measurable, Achievable, Replicable, Time-bound) criteria [8, see
134 Table 3 therein] focus on the decision-makers' receptiveness to suggested indicators during the selection, paving
135 a way to communicate network information with stakeholders and embed network indicators in ecological
136 monitoring and ecosystem health assessments. Network indicators will then need to be evaluated in terms of
137 usefulness to achieve conservation goals [as in O'Connor, 2022ⁱ] and decision-maker receptiveness [as in 8], as
138 we move towards developing ecosystem management and monitoring frameworks that quantitatively and
139 explicitly embed network indicators (see example in Box 2).

140 Second, network ecologists have the opportunity to expand their focus from the development of mathematical
141 tools, theory and theoretical validation to involving decision-makers and meeting their needs [67]. Consensus
142 for conservation goals can be achieved through mixed methodology such as iterative and anonymous Delphi
143 panels [see 68 for applications in ecology]. Engaging stakeholders in this way would ultimately provide
144 valuable guidance to prioritise new fundamental research questions and methodological development. Although
145 they do not ultimately make the decisions, network ecologists must be proactive in this process, especially given
146 the limited time and staffing resources across many institutions where decisions are made. This process takes
147 time and co-production effort, and needs to be initiated by academics who can guide and support practitioners
148 in designing management strategies and making conservation decisions using network information. Academics
149 place a strong focus on the development of tools and knowledge, but ensuring their adoption (particularly for
150 non-academics) will require delivering them in a form that can instantly be used with minimal additional work
151 [69].

152 Finally, network ecologists can take concrete steps to ensure that network-based measures are perceived as
153 relevant by decision-makers. Workshops and stakeholder involvement are essential to bridge the gap between
154 science and practice [69] and can facilitate choosing appropriate metrics [8]. Involving a wide-range of
155 ecosystem-management players, and creating new opportunities to actively involve stakeholders in deciding
156 how network information can be applied, will be key to ensure receptiveness and a speedy uptake of indicators
157 for management planning and actions. Forecasting changes in network structure under environmental and
158 management scenarios (Box 3) and linking network indicators to ecosystem services [17] can enhance
159 receptiveness. This will provide essential information on risks, on boundaries of change given environmental

Box 2 - Assessing the relevance of a potential network indicator for decision-making

Network metrics should be evaluated using criteria important to decision-makers to ensure their relevance as indicators and encourage adoption. In addition to the ROARS and SMART criteria, Fath et al. [8] suggest that effective indicators should also “*describ[e] directional change [of ecosystems], [be] easily communicable to managers and policy makers, [be] integrative and indicative to a known response to a disturbance*” [as per 71], and provide insight to ecosystem functioning and services.

As an example, trophic network robustness to targeted extinctions meets these criteria (Tbl. 1) and can be a useful indicator of ecosystem integrity and stability to environmental change. The structural stability of trophic networks is closely linked to the stability of ecosystem functioning [see review by 72], with trophic interactions considered as ecosystem functions and services (e.g., top-down pest control by predators). Here we show a formulation of robustness derived from earlier works [73–75] that reflects the capacity of a network (or the ecosystem it represents) to withstand cascading extinctions:

$$\text{Robustness} = 1 - \frac{\text{no. secondary extinctions}}{\text{initial no. secondary consumers}}$$

where secondary extinctions are extinctions due to the loss of other species and secondary consumers are consumers of basal species (measured as network species richness minus the number of basal species).

Robustness is easy to interpret (see Specific in Tbl. 1) and to calculate using binary trophic networks, which are more commonly available and can be derived from existing trophic metawebs – this allows us to derive initial (even if coarse) estimates of robustness at large, regional and local scales (see references in Tbl. 1). It also relates to ecological issues that have a firm place in ecosystem management and conservation, and resonate with decision-makers – numerous directives, policies and management frameworks focus on avoiding species extinctions (see examples in Tbl. 1).

Tbl. 1 illustrates the potential of robustness as a network indicator and the process of detailing how it meets the criteria mentioned previously. Evaluating network metrics in this way is crucial for making them more relevant and acceptable to decision-makers, as it demonstrates why and how the indicator can be used effectively.

Table 1: **Relevance of robustness as an indicator.** Dale & Beyler’s [71], ROARS and SMART criteria for good ecological network indicators, as described by Fath et al. [8], and how they apply to robustness of trophic (non-energy flow) networks.

	Description	
Criteria	[as in 8]	How it applies to robustness
Dale & Beyler’s [71]	Describe directional change	Robustness measures loss of species with respect to a given (pre-disturbance) species composition.
	Easily communicable to managers and policy makers	The relationship between robustness and species extinctions is intuitive and easy to understand. See also entry for “Relevant” below.
	Integrative and indicative to a known response to a disturbance	Trophic networks summarise the energy flows in an ecosystem; their structural stability is linked to stability of ecosystem functioning [72]. Robustness measures trophic network responses to disturbances that lead to cascading species extinctions.
ROARS		
Relevant	It relates to an important part of an objective or output	Preventing species extinctions is at the heart of numerous conservation policies, directives and frameworks [e.g., 76,77–79].

Objective	Based on facts, rather than feelings or impressions and thus measurable	Robustness is based on assessments of species composition pre- and post- disturbance.
Available	Data should be readily available or reasonably measurable	At the regional scales, available metawebs [e.g., 43,57] can be combined with species range data (e.g., IUCN ⁱⁱ and GBIF ⁱⁱⁱ) and scenarios of change to assess robustness (see Box 3). Sub-regional/local scale assessments are possible in locations with monitoring data [e.g., 41,42].
Realistic	It should not be too difficult or too expensive to collect the information	Marine and freshwater network data are already being collected as part of monitoring programs and fisheries activities; Terrestrial metawebs exist [43] or can be inferred [80] Methodology to calculate robustness is not overly complex and can be pipelined (see example below).
Specific	The measured changes should be expressed in precise terms	Robustness is calculated as 1 minus the ratio of secondary extinctions to the initial number of secondary consumers. It is scaled from 0-1, with 1 indicating maximum robustness (no secondary extinctions) and 0 indicating no robustness (all secondary consumers went secondarily extinct due to loss of feeding resources).

SMART

Specific	<p>Measured changes should be expressed in precise terms and suggest the direction of actions</p>	<p>See entry for “Specific” above.</p> <p>Maps of robustness can indicate hotspots and priority areas for conservation.</p> <p>Networks with high robustness will indicate ecosystems whose structure is more stable and that could be managed as “safety nets” and/or with more liberal use. Those with low robustness should be further assessed for their uniqueness (e.g., uniqueness of species composition and interactions, of habitat type, etc.) to plan conservation actions.</p>
Measurable	<p>Indicators should be related to things that can be measured in an unambiguous way</p>	<p>In an empirical setting, there may be ambiguity in determining whether an extinction was secondary (due to loss of other species in the network) or primary (due to, e.g., loss of climate suitability).</p> <p>In a modelling setting secondary and primary extinctions can be determined. Null models can be used to test whether forecasted extinctions significantly deviate from random.</p> <p>Uncertainty in both network species composition and structure will need to be recognised and accounted for explicitly whenever possible [e.g., 49]</p>
Achievable	<p>Indicators should be reasonable and possible to reach, and therefore sensitive to changes</p>	<p>See entry for “Available” above.</p> <p>Backcasting and historical observational data can be used to gauge the sensitivity of robustness to past environmental change.</p> <p>Forecasting data can be used to assess robustness boundaries to expected changes and complemented with monitoring data to verify how networks are responding to change.</p>

Replicable	Measurements should be the same when made by different people using the same method	Transparent and freely accessible pipelines can be developed and automated to ensure repeatability.
Time-bound	There should be a time limit within which changes are expected and measured	This likely depends on the species and type of environmental changes considered, given different life cycle histories and species' sensitivities to change.

165

Box 3 - An accessible workflow applying robustness to inform decision-making

Effective decision-making requires indicators based on accessible and reproducible analysis workflows that evaluate a range of scenarios. We demonstrate the potential of robustness with a workflow that uses different network disturbance scenarios and open-access data (Fig. 1). By using extreme scenarios, we can explore the boundaries of robustness to forecasted environmental change. The framework can be applied spatially to identify target areas for management and conservation action (Fig. 2) or to single networks.

Workflow steps:

1. Build local 'reference networks' by combining a regional metaweb of interactions with 'reference' local species presence/absence information ('baseline' referring to any reference period) – species that interact in the metaweb and are locally present, will appear and interact in the local network;
2. For each reference network, calculate the number of secondary consumers (consumers of basal species) and other relevant network metrics (e.g., species and average trophic level, connectance, etc.)
3. Build local 'disturbed networks', by combining the regional metaweb with species ranges projected

166

under different scenarios;

4. Calculate and map robustness and other network metrics (Fig. 2).




Figure 1: **Workflow to calculate robustness.** Simple network metrics like robustness can be incorporated into workflows to assess potential ecosystem fragility to scenarios of disturbance and inform management and decision-making at large scales. See supplemental information online for full workflow details.

Our example explores the lower boundaries of pan-European trophic network robustness by submitting vertebrate networks to two extreme local scenarios: worst-case climate change (CMIP5 RCP 8.5, equivalent to CMIP6 SSP5-8.5), and failure to protect endangered species (IUCN levels: critically endangered, CR, endangered, EN, and vulnerable, VU; Fig. 2). Further analyses could be focused on investigating which species are forecasted to be lost, their roles in the networks and best strategies to protect these networks from a multispecies perspective. For instance, inspecting initial species richness and trophic positions of extinct species can help identify network- and species-level attributes that may be related to robustness (Fig. 2, lower panels). Antunes et al. [17] proposed a similar workflow to calculate network-provided Nature’s contributions to people. Ours differs from theirs in that it requires less sophisticated and less

data-hungry methodological approaches. Together with the accessible automated pipeline [81], this should facilitate and accelerate uptake by practitioners, managers and decision-makers.




Figure 2: Robustness of European vertebrate networks to disturbance scenarios. Extreme scenarios of climate change and of species extinctions can be used to explore (lower) boundaries of network robustness and identify areas where we may expect a high number of cascading (secondary) extinctions and, consequently, larger disruptions to ecosystem functioning and services (upper panels). Further analyses of initial network metrics allow a deeper look into what may drive network robustness by comparing trophic information between primary and secondary extinctions (lower panels, here grouped by quantiles of robustness values). In this example, most networks are very robust to extinctions driven by a) climate change or b) the removal of endangered species listed in IUCN, but several networks in Northern Europe show lower robustness to targeted IUCN extinctions (upper panels). For networks that suffered secondary extinctions (where Robustness < 1; ‘Sec.’ bands on lower panels), larger networks (higher initial species richness) were more robust and, as expected, secondarily extinct species occupied higher trophic positions than primarily extinct species (‘Pri.’ bands). See supplemental information online for more detail. Data and analyses for this figure were adapted from Ceres Barros, PhD thesis, Université Grenoble Alpes, 2017^{iv}.

168

169 Concluding remarks

170 Ecological networks already can and should be used as indicators in biodiversity conservation and ecosystem
 171 management. Sufficient data is available for initial assessments of network structures and responses to change.

172 Additionally, we have relevant network indicators for ecosystem management and conservation that can be
173 weaved into management frameworks and monitoring programs. Starting now ensures that future data will be
174 useful to detect network changes and to address current knowledge gaps.

175 We recognize that the lack of empirical support for theory and scenarios of network responses (including
176 robustness) to environmental change can refrain academics from providing guidance to practitioners.
177 Robustness and extinction studies usually rely on simulations to investigate effects of species losses (rather than
178 observations or experimental removals) and predictions remain mostly untested in the field [82, see Table 1
179 therein for some empirical validation examples]. Overcoming this barrier will require setting up empirical
180 programs that go beyond documenting networks, and towards field and lab studies of network responses to
181 realistic disturbances. Yet, despite this and other limitations (i.e., data, uncertainty, and interpretability
182 challenges), we believe the field is sufficiently mature to make recommendations for ecosystem management
183 and conservation as these programs are implemented.

184 We envision five important aspects for future directions (see also [Outstanding Questions](#)). First, there should be
185 developments addressing evaluation, propagation, and communication of uncertainty in network structure and
186 metrics. It will be key to a) integrate uncertainty robustly into management frameworks and move towards more
187 transparent and informed decisions, but also to b) use existing tools and data to compare known network and
188 ecosystem changes with predictions (e.g. backcasting), estimate boundaries of future network changes
189 (e.g. forecasting), and assess the usefulness of network metrics as indicators of future change. Second, network
190 considerations will need to be explicit in future sampling and monitoring designs, and in ecosystem conservation
191 regulations and decisions. Third, current data, network models and indicators need to be more widely assessed
192 for their usefulness for ecosystem management, which should actively involve stakeholders. Fourth, empirical
193 programs focused on testing and measuring network (metrics') responses to change will need to be set up.
194 Finally, incorporating network information explicitly into conservation will require developing network-based
195 targets—specific, quantified metrics to obtain and thresholds to respect based on whole network characteristics.

Outstanding questions

- How variable is network structure across space and time and does it influence the usefulness of network metrics as indicators of ecosystem functioning and stability?
- What network metrics are ubiquitous, reliable and applicable indicators of ecosystem functioning and stability?

196

- How much can we expect networks to change given uncertainty in future environmental conditions?
- How can current and future monitoring programs be improved to sample network information relevant for management?
- How can we put in place a strong empirical program to validate network indicators, which for now heavily rely on simulations?

197

198 **Acknowledgements**

199 G.D. is funded by the NSERC Postgraduate Scholarship – Doctoral (grant ES D – 558643), the FRQNT
200 doctoral scholarship (grant no. 301750), and the NSERC CREATE BIOS2 program. T.P. is funded by the
201 Wellcome Trust (223764/Z/21/Z), NSERC through the Discovery Grant and Discovery Accelerator
202 Supplements programs, and the Courtois Foundation. WT, LOC and LM acknowledge support from the
203 European Union’s Horizon Europe under grant agreement number 101060429 (project NaturaConnect). JMM
204 acknowledges the support of Horizon Europe project BIOcean5D (award number 101059915) and the French
205 Agence Nationale de la Recherche through LabEx TULIP (ANR-10-LABX-41).

206 **Resources**

207 ⁱ <https://theses.hal.science/tel-04077711>

208 ⁱⁱ <https://www.iucnredlist.org/resources/spatial-data-download>

209 ⁱⁱⁱ <https://www.gbif.org/what-is-gbif>

210 ^{iv} <https://theses.hal.science/tel-01685584>

211 **References**

- 212 1. Franklin, J.F. (1993) [Preserving Biodiversity: Species, Ecosystems, or Landscapes?](#) *Ecological Applications* 3,
202–205
- 213 2. Simberloff, D. (1998) [Flagships, umbrellas, and keystones: Is single-species management passé in the landscape era?](#) *Biological Conservation* 83, 247–257

- 214 3. Heinen, J.H. *et al.* (2020) [Conservation of species interactions to achieve self-sustaining ecosystems](#). *Ecography* 43, 1603–1611
- 215 4. Soulé, M.E. *et al.* (2005) [Strongly Interacting Species: Conservation Policy, Management, and Ethics](#). *BioScience* 55, 168–176
- 216 5. Harvey, E. *et al.* (2017) [Bridging ecology and conservation: From ecological networks to ecosystem function](#). *Journal of Applied Ecology* 54, 371–379
- 217 6. McDonald-Madden, E. *et al.* (2016) [Using food-web theory to conserve ecosystems](#). *Nature Communications* 7, 10245
- 218 7. Tylianakis, J.M. *et al.* (2010) [Conservation of species interaction networks](#). *Biological Conservation* 143, 2270–2279
- 219 8. Fath, B.D. *et al.* (2019) [Ecological network analysis metrics: The need for an entire ecosystem approach in management and policy](#). *Ocean & Coastal Management* 174, 1–14
- 220 9. Leffland, K. *et al.* (1998) *Comparing environmental impact data on cleaner technologies*, Office for official publications of the European communities
- 221 10. Gregr, E.J. *et al.* (2020) [Cascading social-ecological costs and benefits triggered by a recovering keystone predator](#). *Science* 368, 1243–1247
- 222 11. Geary, W.L. *et al.* (2020) [A guide to ecosystem models and their environmental applications](#). *Nature Ecology & Evolution* 4, 1459–1471
- 223 12. Rapacciuolo, G. (2019) [Strengthening the contribution of macroecological models to conservation practice](#). *Global Ecology and Biogeography* 28, 54–60
- 224 13. Executive Secretary of the Convention on Biological Diversity (2022) *Expert input to the Post-2020 Global Biodiversity Framework: Transformative actions on all drivers of biodiversity loss are urgently required to achieve the global goals by 2050*
- 225 14. Leadley, P. *et al.* (2022) [Achieving global biodiversity goals by 2050 requires urgent and integrated actions](#). *One Earth* 5, 597–603
- 226 15. Araújo, M.B. and Alagador, D. (2024) [Expanding European protected areas through rewilding](#). *Current Biology* 34, 3931–3940.e5
- 227 16. Gaüzère, P. *et al.* (2023) [Dissimilarity of vertebrate trophic interactions reveals spatial uniqueness but functional redundancy across Europe](#). *Current Biology* 33, 5263–5271.e3
- 228 17. Antunes, A.C. *et al.* (2024) [Linking biodiversity, ecosystem function, and Nature’s contributions to people: A macroecological energy flux perspective](#). *Trends in Ecology & Evolution* 39, 427–434
- 229 18. CBD (2022) *Decision adopted by the conference of the parties to the convention on biological diversity 15/4. Kunming-montreal global biodiversity framework*
- 230 19. Keyes, A.A. *et al.* (2021) [An ecological network approach to predict ecosystem service vulnerability to species losses](#). *Nature Communications* 12, 1586

- 231 20. Poisot, T. *et al.* (2016) [The structure of probabilistic networks](#). *Methods in Ecology and Evolution* 7, 303–312
- 232 21. Mills, L.S. *et al.* (1993) [The Keystone-Species Concept in Ecology and Conservation](#). *BioScience* 43, 219–224
- 233 22. CBD (2022) Saving endangered keystone species: Key to ecosystem restoration *Convention on Biological Diversity* <https://www.cbd.int/article/saving-endangered-keystone-species>
- 234 23. Europe, R. (2019) The Keystone Concept. *Rewilding Europe*
- 235 24. Jordán, F. (2009) [Keystone species and food webs](#). *Philosophical Transactions of the Royal Society B: Biological Sciences* 364, 1733–1741
- 236 25. Shukla, I. *et al.* (2023) [The diversity of animals identified as keystone species](#). *Ecology and Evolution* 13, e10561
- 237 26. Ripple, W.J. *et al.* (2014) [Status and Ecological Effects of the World’s Largest Carnivores](#). *Science* 343, 1241484
- 238 27. Kotliar, N.B. *et al.* (1999) [A Critical Review of Assumptions About the Prairie Dog as a Keystone Species](#). *Environmental Management* 24, 177–192
- 239 28. Miller, B. *et al.* (2000) [The Role of Prairie Dogs as a Keystone Species: Response to Stapp](#). *Conservation Biology* 14, 318–321
- 240 29. Agency, P.C. (2021) *Recovery Strategy and Action Plan for the Black-tailed Prairie Dog (Cynomys ludovicianus) in Canada*, Parks Canada Agency
- 241 30. Gouveia, C. *et al.* (2021) [Combining centrality indices: Maximizing the predictability of keystone species in food webs](#). *Ecological Indicators* 126, 107617
- 242 31. Gonzalez, A. *et al.* (2017) [Spatial ecological networks: Planning for sustainability in the long-term](#). *Current Opinion in Environmental Sustainability* 29, 187–197
- 243 32. Tonn, B. *et al.* (2000) [A Framework for Understanding and Improving Environmental Decision Making](#). *Journal of Environmental Planning and Management* 43, 163–183
- 244 33. Hortal, J. *et al.* (2015) [Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity](#). *Annual Review of Ecology, Evolution, and Systematics* 46, 523–549
- 245 34. Jordano, P. (2016) [Chasing Ecological Interactions](#). *PLOS Biology* 14, e1002559
- 246 35. Mestre, F. *et al.* (2022) [Disentangling food-web environment relationships: A review with guidelines](#). *Basic and Applied Ecology* 61, 102–115
- 247 36. Poisot, T. *et al.* (2021) [Global knowledge gaps in species interaction networks data](#). *Journal of Biogeography* 48, 1552–1563
- 248 37. Windsor, F.M. *et al.* (2023) [Using ecological networks to answer questions in global biogeography and ecology](#). *Journal of Biogeography* 50, 57–69
- 249 38. Jordano, P. (2016) [Sampling networks of ecological interactions](#). *Functional Ecology* 30, 1883–1893

- 250 39. Vázquez, D. *et al.* (2022) [Ecological interaction networks. What we know, what we don't, and why it matters.](#) *Ecologia Austral* 32, 670–697
- 251 40. Cameron, E.K. *et al.* (2019) [Uneven global distribution of food web studies under climate change.](#) *Ecosphere* 10, e02645
- 252 41. Bonnaffé, W. *et al.* (2021) [Comparison of size-structured and species-level trophic networks reveals antagonistic effects of temperature on vertical trophic diversity at the population and species level.](#) *Oikos* 130, 1297–1309
- 253 42. Danet, A. *et al.* (2021) [Species richness and food-web structure jointly drive community biomass and its temporal stability in fish communities.](#) *Ecology Letters* 24, 2364–2377
- 254 43. Maiorano, L. *et al.* (2020) [TETRA-EU 1.0: A species-level trophic metaweb of European tetrapods.](#) *Global Ecology and Biogeography* 29, 1452–1457
- 255 44. McLeod, A. *et al.* (2021) [Sampling and asymptotic network properties of spatial multi-trophic networks.](#) *Oikos* 130, 2250–2259
- 256 45. Strydom, T. *et al.* (2023) [Graph embedding and transfer learning can help predict potential species interaction networks despite data limitations.](#) *Methods in Ecology and Evolution* 14, 2917–2930
- 257 46. Albouy, C. *et al.* (2019) [The marine fish food web is globally connected.](#) *Nature Ecology & Evolution* 3, 1153–1161
- 258 47. Braga, J. *et al.* (2019) [Spatial analyses of multi-trophic terrestrial vertebrate assemblages in Europe.](#) *Global Ecology and Biogeography* 28, 1636–1648
- 259 48. Galiana, N. *et al.* (2021) [The spatial scaling of food web structure across European biogeographical regions.](#) *Ecography* 44, 653–664
- 260 49. Dansereau, G. *et al.* (2024) [Spatially explicit predictions of food web structure from regional-level data.](#) *Philosophical Transactions of the Royal Society B: Biological Sciences* 379, 20230166
- 261 50. Johnson, S. *et al.* (2023) [Field validation as a tool for mitigating uncertainty in species distribution modeling for conservation planning.](#) *Conservation Science and Practice* 5, e12978
- 262 51. Morales-Castilla, I. *et al.* (2015) [Inferring biotic interactions from proxies.](#) *Trends in Ecology & Evolution* 30, 347–356
- 263 52. Adams, M.P. *et al.* (2020) [Informing management decisions for ecological networks, using dynamic models calibrated to noisy time-series data.](#) *Ecology Letters* 23, 607–619
- 264 53. Shirey, V. and Rabinovich, J. (2024) [Climate change-induced degradation of expert range maps drawn for kissing bugs \(Hemiptera: Reduviidae\) and long-standing current and future sampling gaps across the Americas.](#) *Memórias do Instituto Oswaldo Cruz* 119, e230100
- 265 54. Van Kleunen, L.B. *et al.* (2023) [Decision-making under uncertainty for species introductions into ecological networks.](#) *Ecology Letters* 26, 983–1004
- 266 55. Pollock, L.J. *et al.* (2020) [Protecting Biodiversity \(in All Its Complexity\): New Models and Methods.](#) *Trends in Ecology & Evolution* 35, 1119–1128

- 267 56. Thuiller, W. *et al.* (2019) [Uncertainty in ensembles of global biodiversity scenarios](#). *Nature Communications* 10, 1446
- 268 57. Frelat, R. *et al.* (2022) [Food web structure and community composition: A comparison across space and time in the North Sea](#). *Ecography* 2022
- 269 58. Kortsch, S. *et al.* (2015) [Climate change alters the structure of arctic marine food webs due to poleward shifts of boreal generalists](#). *Proceedings of the Royal Society B: Biological Sciences* 282, 20151546
- 270 59. Kortsch, S. *et al.* (2019) [Food-web structure varies along environmental gradients in a high-latitude marine ecosystem](#). *Ecography* 42, 295–308
- 271 60. Trifonova, N. *et al.* (2015) [Spatio-temporal Bayesian network models with latent variables for revealing trophic dynamics and functional networks in fisheries ecology](#). *Ecological Informatics* 30, 142–158
- 272 61. Brunner, S.H. *et al.* (2016) [A backcasting approach for matching regional ecosystem services supply and demand](#). *Environmental Modelling & Software* 75, 439–458
- 273 62. O'Connor, L.M.J. *et al.* (2024) [Vulnerability of terrestrial vertebrate food webs to anthropogenic threats in Europe](#). *Global Change Biology* 30, e17253
- 274 63. Cirtwill, A.R. *et al.* (2024) [Species motif participation provides unique information about species risk of extinction](#). *Journal of Animal Ecology* 93, 731–742
- 275 64. Stouffer, D.B. and Bascompte, J. (2010) [Understanding food-web persistence from local to global scales](#). *Ecology Letters* 13, 154–161
- 276 65. Baiser, B. *et al.* (2010) [Connectance determines invasion success via trophic interactions in model food webs](#). *Oikos* 119, 1970–1976
- 277 66. Romanuk, T.N. *et al.* (2017) [Chapter Five - Robustness Trade-Offs in Model Food Webs: Invasion Probability Decreases While Invasion Consequences Increase With Connectance](#). In *Advances in Ecological Research* 56 (Bohan, D. A. *et al.*, eds), pp. 263–291, Academic Press
- 278 67. Bodner, K. *et al.* (2021) [Bridging the divide between ecological forecasts and environmental decision making](#). *Ecosphere* 12, e03869
- 279 68. Mukherjee, N. *et al.* (2015) [The Delphi technique in ecology and biological conservation: Applications and guidelines](#). *Methods in Ecology and Evolution* 6, 1097–1109
- 280 69. Jung, M. *et al.* (2024) [An assessment of the state of conservation planning in Europe](#). *Philosophical Transactions of the Royal Society B: Biological Sciences* 379, 20230015
- 281 70. Polasky, S. *et al.* (2011) [Decision-making under great uncertainty: Environmental management in an era of global change](#). *Trends in Ecology & Evolution* 26, 398–404
- 282 71. Dale, V.H. and Beyeler, S.C. (2001) [Challenges in the development and use of ecological indicators](#). *Ecological Indicators* 1, 3–10
- 283 72. Saint-Béat, B. *et al.* (2015) [Trophic networks: How do theories link ecosystem structure and functioning to stability properties? A review](#). *Ecological Indicators* 52, 458–471

- 284 73. Dunne, J.A. *et al.* (2002) [Network structure and biodiversity loss in food webs: Robustness increases with connectance](#). *Ecology Letters* 5, 558–567
- 285 74. Dunne, J.A. *et al.* (2004) [Network structure and robustness of marine food webs](#). *Marine Ecology Progress Series* 273, 291–302
- 286 75. Solé, R.V. and Montoya, M. (2001) [Complexity and fragility in ecological networks](#). *Proceedings of the Royal Society of London. Series B: Biological Sciences* 268, 2039–2045
- 287 76. Directive, B. (2009) Directive 2009/147/EC of the European Parliament and of the Council of 30 November 2009 on the conservation of wild birds
- 288 77. Law, C.W.P. (2018) Wildlife Protection Law of the People’s Republic of China
- 289 78. Amendment Bill, W.L. (Protection) (2021) Wild Life (Protection) Amendment Bill
- 290 79. UN Convention on Biological Diversity (1992) UN Convention on Biological Diversity
- 291 80. Strydom, T. *et al.* (2022) [Food web reconstruction through phylogenetic transfer of low-rank network representation](#). *Methods in Ecology and Evolution* 13, 2838–2849
- 292 81. Barros, C. (2024) [CeresBarros/TrophicNetRobWF: V0.0.0.9000](#)Zenodo
- 293 82. Morton, D.N. *et al.* (2022) [Merging theory and experiments to predict and understand coextinctions](#). *Trends in Ecology & Evolution* 37, 886–898